

- of fresh-cut 'Kent' mango (*Mangifera indica* L.). Post Biol Technol 2009; 51(3): 384-90.
<http://dx.doi.org/10.1016/j.postharvbio.2008.09.003>
- [65] Tezcan F, Gultekin-Ozguven M, Diken T, Ozcelik B, Bedia Erim F. Antioxidant activity and total phenolic, organic acid and sugar content in commercial pomegranate juices. Food Chem 2009; 115: 873-77.
<http://dx.doi.org/10.1016/j.foodchem.2008.12.103>
- [66] Bail S, Stuebiger G, Krist S, Unterweger H. Characterisation of various grape seed oils by volatile compounds, triacylglycerol composition, total phenols and antioxidant capacity. Food Chem 2008; 108(3): 1122-32.
<http://dx.doi.org/10.1016/j.foodchem.2007.11.063>
- [67] Usenik V, Fabcic J, Stampar F. Sugars, organic acids, phenolic composition and antioxidant activity of sweet cherry (*Prunus avium* L.). Food Chem 2008; 107: 185-92.
<http://dx.doi.org/10.1016/j.foodchem.2007.08.004>
- [68] Odriozola-Serrano I, Soliva-Fortuny R, Martin-Belloso O. Effect of minimal processing on bioactive compounds and colour attributes of fresh-cut tomatoes. LWT 2008; 41: 217-26.
<http://dx.doi.org/10.1016/j.lwt.2007.03.002>
- [69] Netzel M, Netzel G, Tian Q, Schwartz S, Konczak I. Native Australian fruits – a novel source of antioxidants for food. Innov Food Sci Emerg Technol 2007; 8: 339-46.
<http://dx.doi.org/10.1016/j.ifset.2007.03.007>
- [70] Park YS, Jung ST, Kang SG, Delgado-Licon E, Leticia Martinez Ayala A, et al. Drying of persimons (*Diospyros kaki* L.) and the following changes in the studied bioactive compounds and total radical scavenging activities. LWT 2006; 39: 748-55.
<http://dx.doi.org/10.1016/j.lwt.2005.05.014>
- [71] Assimopoulou AN, Zlatanov SN, Papageorgiou VP. Antioxidant activity of natural resins and bioactive triterpenes in oil substrates. Food Chem 2005; 92: 721-27.
<http://dx.doi.org/10.1016/j.foodchem.2004.08.033>
- [72] Ayoughi F, Barzegar M, Sahari MA, Naghdi Badi HA. Chemical compositions of essential oils of *Artemisia dracunculoides* L. and endemic *Matricaria chamomilla* L. and an evaluation of their antioxidative effects. J Agric Sci Technol 2011; 13: 79-88.
- [73] Darughe F, Barzegar M, Sahari MA. Antioxidant and antifungal activity of Coriander (*Coriandrum sativum* L.) essential oil in cake. Inter Food Res J 2012; 19(3): 1253-60.
- [74] Fathi A, Sahari MA, Zangiabadi M, Barzegar M. Application of *Satureja hortensis* L. and *Zataria multiflora* Boiss. essential oils as two natural antioxidants in soybean oil during microwave heating. J Med Plants 2011; 10: 12-21.
- [75] Fazel M, Sahari MA, Barzegar M. Determination of main tea seed oil antioxidants and their effects on common kilka oil. Inter Food Res J 2008; 15 (2): 209-17.
- [76] Fazel M, Sahari MA, Barzegar M. Comparison of tea and sesame seed oils as two natural antioxidants in a fish oil model system by radical scavenging activity. Inter J Food Sci Nutr 2009; 60(7): 567-76.
<http://dx.doi.org/10.3109/09637480801987625>
- [77] Hassan Zadeh A, Sahari MA, Barzegar M. Optimization of the v-3 extraction as a functional food from flaxseed. Inter J Food Sci Nutr 2008; 59(6): 526-34.
<http://dx.doi.org/10.1080/09637480701565935>
- [78] Khaki M, Sahari MA, Barzegar M. Evaluation of antioxidant and antimicrobial effects of chamomile (*Matricaria chamomilla* L.) essential oil on cake shelf life. J Med Plants 2012; 11: 9-18.
- [79] Noorolahi Z, Sahari MA, Barzegar M, Naghdi Badi HA. Evaluation Antioxidant and Antimicrobial Effects of Cinnamon Essential Oil and Echinacea Extract in Kolompe. J Med Plants 2012; 11: (In Press).
- [80] Rajaei A, Barzegar M, Sahari M. A. Comparison of antioxidative effect of tea and sesame seed oils extracted by different methods. J Agric Sci Technol 2008; 10: 345-50.
- [81] Rajaei A, Barzegar M, Mohabati Mobarez A, Sahari MA, Hamidi Esfahani, Z. Antioxidant, anti-microbial and antimutagenicity activities of pistachio (*Pistachia vera*) green hull extract. Food Chem Toxicol 2010; 48: 107-12.
<http://dx.doi.org/10.1016/j.foodtox.2009.09.023>
- [82] Sabouri Z, Barzegar M, Sahari MA, Naghdi Badi HA. Antioxidant and antimicrobial potential of *Echinacea purpurea* extract and its effect on extension of cake shelf life. J Med Plants 2012; 11: 28-40.
- [83] Soleiman H, Barzegar M, Sahari MA, Naghdi Badi HA. An investigation on the antioxidant activities of *Hyssopus officinalis* L. and *Echinacea purpurea* L. plant extracts in oil model system. J Med Plants 2011; 10: 61-72.
- [84] Taha Nejad M, Barzegar M, Sahari MA, Naghdi Badi HA. Evaluation of antioxidant activity of *Lavandula angustifolia* essential oil in crude soybean oil system. J Med Plants 2011; 11(1): 127-140 [In Farsi].
- [85] Taha Nejad M, Barzegar M, Sahari MA, Naghdi Badi HA. Evaluation of antioxidant activity of *Malva sylvestris* L. extract in crude soybean oil system. J Med Plants 2012; 11(2): 86-97 [In Farsi].
- [86] Vahidian H, Sahari MA, Barzegar M, Naghdi Badi HA. Application of *Zataria multiflora* Boiss. and *Satureja hortensis* L. essential oils as two natural antioxidants in mayonnaise formulated with linseed oil. J Med Plants 2012; 11: 69-79.
- [87] Yasoubi P, Barzegar M, Sahari MA, Azizi MA. Total phenolic contents and antioxidant activity of pomegranate (*Punica granatum* L.) peel extracts. J Agric Sci Technol 2007; 9: 35-42.
- [88] Zangiabadi M, Sahari MA, Barzegar M, Naghdi Badi HA. *Zataria multiflora* and *Bunium persicum* essential oils as two natural antioxidants. J Med Plants 2012; 11: 8-21.